

AN ATLAS OF PHANEROZOIC CLADE DIVERSITY DIAGRAMS

J. JOHN SEPKOSKI, Jr. and MICHAEL L. HULVER

Department of the Geophysical Sciences, University of Chicago

Clade diversity diagrams are spindle-shaped graphs that summarize patterns of taxonomic evolution within higher taxa through geologic time. Most clade diversity diagrams are constructed about a central axis that represents time (scaled either metrically or ordinally, by stratigraphic interval). Some measure or estimate of taxonomic diversity (or “richness”) is then plotted symmetrically about the axis to give the diagram an overall spindle shape (e.g., Figure 1).

Diversity diagrams for individual clades convey information about their size, shape, and variability in the fossil record (cf. Gould et al., 1977). Such “morphologic” information is valuable for assessing how evolutionary rates (that is, rates of origination and extinction) vary within the taxa through geologic time. Clade diversity diagrams for groups of higher taxa hypothesized to be related by phylogeny or by function are useful for comparisons of the histories of the taxa. Common patterns of expansion or contraction may relate to general factors governing all taxa, whereas reciprocal patterns may be interpretable as negative interactions between pairs of ecologically similar taxa (e.g., Simpson, 1953; Bambach, this volume). Sets of clade diversity diagrams also are useful for summarizing variation among large numbers of clades for the purpose of testing general macroevolutionary models (e.g., Raup et al., 1973; Gould et al., 1977).

THE MARINE FOSSIL RECORD




Figure 1. Families within classes of fossil marine mammals.

This chapter presents a collection of clade diversity diagrams which we hope will be useful for examining the general histories of a wide variety of animal taxa. The main body of the chapter is a series of 12 figures displaying spindle diagrams for orders, classes, and phyla of both marine and nonmarine (or "continental") animals for the whole of the Phanerozoic (including the Vendian). Nearly all of the diagrams are plotted at a uniform taxonomic and temporal resolution, specifically that of familial diversity per stratigraphic stage. The taxonomic rank of family is used simply because comprehensive data with good stratigraphic resolution can be obtained for all animal groups at this level. Although families do not display all of the detail of the fossil record, they should be sufficiently sensitive to show major evolutionary trends and patterns with characteristic timescales of fives to tens of million years (see also Sepkoski, 1979, 1982a, Raup and Sepkoski, 1982).

The clade diversity diagrams in most of the figures are formatted in strips that have time in the vertical dimension. Most of the strips are scaled from 625 myr at their bottoms to approximately 1 myr BP at their tops. (No data on Recent diversity are directly included in the diagrams.) Geologic eras and systems are indicated at the lefthand ends of the strips, with eras denoted by *Cz* = Cenozoic, *Mz* = Mesozoic, *Pz* = Paleozoic, and *pe* = latest Precambrian; systems are denoted by standard symbols, with *V* = Vendian. The widths of the clade diversity diagrams in each strip indicate the numbers of families known from direct fossil evidence or from interpolation between known occurrences to be present in the "clades" in each of 80 stratigraphic stages or comparable intervals (see Table 1 in Sepkoski, 1982b for a listing of the stages used). A scale for the familial diversities appears in the lower righthand part of most of the figures. All of the diagrams were produced with an IBM Personal Computer and Epson dot-matrix printer.

The first two figures in this chapter contain class-level summaries of the entire Phanerozoic fossil record. Figure 1 displays clade diversity diagrams for the 87 classes and 15 unique, problematic genera that have representatives in the marine fossil record. This illustration is an updated version of Figure 1 in Sepkoski (1981) with corrections based on new data in Sepkoski (1982b). The second figure in this chapter summarizes the continental fossil record. The diversity diagrams display numbers of freshwater and terrestrial families within the 39 animal classes known from the nonmarine fossil record; data on the classes were compiled from the literature sources listed in Table 1. Also shown at the bottom of Figure 2 are clade diversity diagrams for numbers of *species* within the 13 taxonomic divisions of the tracheophytes and bryophytes; the data for these diagrams were taken from Niklas, Tiffney and Knoll (this volume).

The next eight figures illustrate a breakdown of the class-level clades into their constituent orders. Time and diversity in all diagrams are plotted at the

TABLE 1. Principal literature sources of information on the taxonomy and stratigraphy of continental animal families.

<i>Taxon</i>	<i>References</i>
MOLLUSCA:	Davies (1971), Henderson (1935), Moore, Teichert, and Robison (1953-1982), Orlov (1958-1964), Solem and Yochelson (1979), Taylor and Sohi (1962).
ARTHROPODA: (excl. insects)	Cooper (1964), Harland et al. (1967), Kukalová-Peck (1973), Moore, Teichert, and Robison (1953-1982), Morris (1979), Mundel (1979), Orlov (1958-1964), Piveteau (1952-1969), Rolfe et al. (1983), Schram (1969), Schram and Schram (1979).
INSECTA:	Barthel (1978), Bode (1953), Burnham (1978), Carpenter (1976, 1979, 1980), Evans (1956), Grande (1980), Harland et al. (1967), Hoganson and Ashworth (1982), Jarzembowski (1980), Kukalová (1966, 1969), Kukalová-Peck (1973, 1975), MacLeod (1970), Müller (1963-1970), Orlov (1958-1964), Piveteau (1952-1969), Rodendorf (1968), Rolfe et al. (1983), Whalley (1980), Wighton (1982), Wilson (1978).
CHORDATA:	Brodkorb (1967, 1971, 1978), Carroll (1977), Charig et al. (1976), Denison (1978, 1979), Eisenberg (1981), Estes (1981), Grande (1980), Harland et al. (1967), Kuhn (1969), Lillegraven et al. (1979), Miynarski (1976), Moy-Thomas and Miles (1971), Nelson (1976), Olsen and Galton (1977), Orlov (1958-1964), Romer (1966), Russell (1975), Steel (1970, 1973), Zangerl (1981).
OTHERS:	Clark (1969), Conway Morris (1981), Conway Morris et al. (1982), Harland et al. (1967), Kukalová-Peck (1973), Moore, Teichert and Robison (1953-1982), Schram (1979), Southcott and Lange (1971), Thompson and Jones (1980).





Figure 2. Families within classes of continental (i.e., terrestrial and freshwater) fossil



animals and species within divisions of continental plants.

THE CONTINENTAL FOSSIL RECORD




Figure 3. Families within orders of fossil "Protozoa," Porifera, and Coelenterata.




Figure 4. Families within orders of fossil Bryozoa, Brachiopoda, Annelida, and Hemichordata.





Figure 5. Families within orders of fossil marine Mollusca



MARINE MOLLUSCS





Figure 6. Families within orders of fossil marine Arthropoda.



MARINE ARTHROPODS





Figure 7. Families within orders of fossil Echinodermata.







Figure 8. Families within orders of marine Vertebrata.



MARINE VERTEBRATES




Figure 9. Families within orders of fossil nonmarine Arthropoda.




Figure 10. Families within orders of fossil nonmarine Vertebrata.

same relative scale as in Figures 1 and 2 in order to facilitate comparison. Figures 3 and 4 display family-level clade diagrams for orders within the moderately diverse marine phyla: the Protozoa, Porifera, Coelenterata, Bryozoa, Brachiopoda, Annelida, and Hemichordata. (The set of clade diversity diagrams for the Annelida includes several taxa of questionable affinities which might best be considered *incertae sedis*; these are in the group of diagrams beginning with Cribricyathida and ending with Cornulitidae.) The more diverse marine phyla are represented in Figures 5 to 8. Figure 5 displays orders of marine molluscs; Figure 6 orders of marine arthropods; Figure 7 orders of echinoderms; and Figure 8 orders of marine vertebrates.

The two large phyla of continental animals, the nonmarine Arthropoda and Chordata, are featured in Figures 9 and 10. Nonmarine taxa have been segregated from their marine relatives because we believe that the land and sea are best treated as separate major arenas of evolution (see also Boucot, 1983). Despite the fact that some continental clades contain secondary species which alternate between marine and nonmarine habitats, and that all clades ultimately had their origins in the oceans, the great majority of continental animals evolved *in situ*, isolated from evolutionary activity in the seas. Thus, the segregation of marine and continental taxa enhances assessment of evolutionary patterns within the two arenas as well as comparisons between them. Note that the time axes for the continental clade diversity diagrams in Figures 9 and 10 have been truncated below the Silurian; this is because there is virtually no nonmarine fossil record prior to the mid Paleozoic (see Boucot and Janis, 1983).

The final pair of figures in this chapter (Figures 11 and 12) contains 14 diversity diagrams for families within entire phyla, split again between marine and continental. These diagrams are formatted somewhat differently than in the preceding figures. The spindles have been cut in half and rotated so that the time axis runs horizontally. This arrangement permits easier assessment of the times and magnitudes of diversity change but impedes comparison of changes between groups.

The use of a single level of taxonomic and stratigraphic resolution in all clade diversity diagrams is intended to aid interpretation and comparison of patterns among the various taxa. However, the constancy of resolution does not imply a uniformity of quality throughout the data. The accuracy of the taxonomic and stratigraphic information varies considerably among the taxonomic groups. In general, the quality is much better for marine taxa than for nonmarine taxa. Also, as should be expected, the fossil data are much better (and much more complete) for heavily skeletonized animals than for soft-bodied and lightly sclerotized animals. In fact, many of the diagrams for the latter groups reflect little more than the geologic distribution of Lagerstätten that preserve unusual fossils. This is particularly evident in the long, thin




Figure 11. Familial diversity of selected marine animal phyla.

clade diagrams for such extant groups as the Nemertinea and Priapulida (Figure 1); these diagrams show only the extension of stratigraphic ranges from the Recent to the one or more Lagerstätten that happen to contain the groups' early members.

Much of the character of the diversity diagrams for some large clades, such as insects (Figures 2 and 9), also represents the effects of Lagerstätten. For the insects, the more important Lagerstätten include the Upper Carboniferous




Figure 12. Familial diversity of selected marine and continental animal phyla.

siderite concretion deposits of North America and Europe, the mid-Permian lake deposits of Kansas and Kazakhstan, the Eocene Green River deposits of Wyoming, and especially the Oligocene Baltic Amber of northern Europe. The Baltic Amber alone contributes most of the Cenozoic bulge in the clade diversity diagrams for both insects and other lightly sclerotized terrestrial arthropods (Figures 2, 9, and 12). The effects of Lagerstätten, or of their non-occurrence, are even seen in some well-skeletonized groups with fairly

extensive fossil records. The drop in the diversity of continental vertebrates in the Jurassic (Figure 12), for example, probably reflects largely a paucity of fossiliferous continental deposits between the Rhaetian and Tithonian (see also Carroll, 1977; Padian and Clemens, 1985, this volume).

These shortcomings of the fossil record, along with the problems associated with family-level data and 5 to 10 myr-long stages, do limit the value of the clade diversity diagrams presented here. However, we believe that a great deal still can be learned from them about the shape of evolution—about the success and failure of taxa and about the apparent order, or disorder, in their radiations and extinctions. Thus, we hope that this “atlas” will aid in the assessment and interpretation of evolutionary history as well as serve as a baseline for the compilation of more accurate and detailed diversity data.

ACKNOWLEDGEMENTS

We thank K. J. Niklas, B. H. Tiffney, and A. H. Knoll for permission to reproduce their data on continental plant diversity in Figure 2. We also thank J. Cracraft, P. Crane, and J. A. Hopson for help and advice during compilation of data on continental animals. Production of this paper received partial support from NSF Grant DEB81-08890 to J. J. S.

REFERENCES

- Bambach, R. K., 1985. Classes and adaptive variety: The ecology of diversification in marine faunas through the Phanerozoic. In Valentine, J. W. (ed.), *Phanerozoic diversity patterns: Profiles in macroevolution*, Princeton, N.J.: Princeton Univ. Press and Amer. Assoc. Adv. Sci. (this volume).
- Barthel, K. W., 1978. *Solnhofen. Ein Blick in die Erdgeschichte*. Thun, Switzerland: Ott Verlag, 393.
- Bode, A., 1953. Die Insektenfauna des Ostniedersächsischen Oberen Lias. *Palaeontographica Abt. A*, 103:1-375.
- Boucot, A. J., 1983. Does evolution take place in an ecological vacuum? II. *J. Paleontol.*, 57:1-30.
- Boucot, A. J., and Janis, C., 1983. Environment of the early Paleozoic vertebrates. *Palaeogeogr., Palaeoclimat., Palaeoecol.* 41:251-288.
- Brodkorb, P., 1967. Catalogue of fossil birds, Part 3 (Ralliformes, Ichthyornithiformes, Charadriiformes). *Florida State Mus. Bull., Biol. Sci.* 11(3): 99-220.
- Brodkorb, P., 1971. Catalogue of fossil birds, Part 4 (Columbiformes through Piciformes). *Florida State Mus. Bull., Biol. Sci.* 15(4):163-266.
- Brodkorb, P., 1978. Catalogue of fossil birds, Part 5 (Passeriformes). *Florida*

- State Mus. Bull., Biol. Sci. 23(3):139-228.
- Burnham, L., 1978. Survey of social insects in the fossil record. *Psyche* 85: 85-133.
- Carpenter, F. M., 1976. The Lower Permian insects of Kansas, Part 12. Prothoptera (continued), Neuroptera, additional Palaeodictyoptera, and families of uncertain position. *Psyche* 83:336-378.
- Carpenter, F. M., 1979. Lower Permian insects from Oklahoma, Part 2. Orders Ephemeroptera and Palaeodictyoptera. *Psyche* 86:261-290.
- Carpenter, F. M., 1980. Studies on North American Carboniferous insects. 6. Upper Carboniferous insects from Pennsylvania. *Psyche* 87:107-119.
- Carroll, R. L., 1977. Patterns of amphibian evolution: An extended example of the incompleteness of the fossil record. In Hallam, A. (ed.), *Patterns of evolution*, Amsterdam: Elsevier, 405-437.
- Charig, A. J., Krebs, B., Sues, H. -D., and Westphal, F., 1976. Thecodontia. *Encyclopedia of paleoherpetology*, Pt. 13. Stuttgart: Gustav Fischer Verlag, 137.
- Clark, R. B., 1969. Systematics and phylogeny: Annelida, Echiura, Sipuncula. In Florkin, M. and Schoer, B. T. (eds.), *Chemical zoology*, v. 4, New York: Academic Press, 1-68.
- Conway Morris, S., 1981. Parasites and the fossil record. *Parasitology* 82:489-509.
- Conway Morris, S., Pickerill, R. K., and Harland, T. L., 1982. A possible annelid from the Trenton Limestone (Ordovician) of Quebec, with a review of fossil oligochaetes and other annulate worms. *Can. J. Earth Sci.* 19: 2150-2157.
- Cooper, K. W., 1964. The first fossil tardigrade: *Beorn leggi* Cooper, from Cretaceous amber. *Psyche* 71:41-48.
- Davies, A. M., 1971. Tertiary faunas. v.1. The composition of Tertiary faunas. Amsterdam: Elsevier, 571.
- Denison, R., 1978. Placodermi. *Handbook of paleoichthyology*, v.2. Stuttgart: Gustav Fischer Verlag, 128.
- Eisenberg, J. F., 1981. The mammalian radiations. Chicago: Univ. Chicago Press, 610.
- Estes, R., 1981. Gymnophiona, Caudata. *Encyclopedia of paleoherpetology*, Pt. 2. Stuttgart: Gustav Fischer Verlag, 115.
- Evans, J. W., 1956. Palaeozoic and Mesozoic Hemiptera (insects). *Aust. Zool. J.* 4:165-258.
- Gould, S. J., Raup, D. M., Sepkoski, J. J. Jr., Schopf, T. J. M., and Simberloff, D. S., 1977. The shape of evolution: A comparison of real and random clades. *Paleobiol.* 3:23-40.
- Grande, L., 1980. Paleontology of the Green River Formation with a review of the fish fauna. *Geol. Surv. Wyoming Bull.* 63.
- Harland, W. B., et al., eds., 1967. *The fossil record*. London: Geol. Soc. London, 828.
- Henderson, J., 1935. Fossil non-marine Mollusca of North America. *Geol. Soc. Amer. Spec. Pap.* 3, 313.
- Hoganson, J. W. and Ashworth, A. C., 1982. The late-glacial climate of the Chilean Lake region implied by fossil beetles. *Third N. Am. Paleont. Conv., Proc.* 1:251-256.
- Jarzembowski, E. A., 1980. Fossil insects from Bembridge Marls, Palaeogene of the Isle of Wight, southern England. *Br. Mus. Nat. Hist. Bull. (Geol.)* 33:237-293.
- Kuhn, O., 1969. Cotylosauria. *Encyclopedia of paleoherpetology*, Pt. 6. Stuttgart: Gustav Fischer Verlag, 89.
- Kukalová, J., 1966. Protelytroptera from the Upper Permian of Australia, with a discussion of Protocoleoptera and Paracoleoptera. *Psyche* 73: 89-111.
- Kukalová, J., 1969. Revisional study of the order Palaeodictyptera in the Upper Carboniferous shales of Commeny, France. Part 1. *Psyche* 76: 163-215.
- Kukalová-Peck, J., 1973. A phylogenetic tree of the animal kingdom (including orders and higher categories). Canada: Ottawa, Nat. Museums of Canada, Publ. in Zool. no. 8, 78.
- Kukalová-Peck, J., 1975. Megasecoptera from the Lower Permian of Moravia. *Psyche* 82:1-19.
- Lillegraven, J. A., Kielan-Jaworowska, Z., and Clemens, W. A., eds., 1979. Mesozoic mammals. Berkeley: Univ. Calif Press, 311.
- MacLeod, E. G., 1970. The Neuroptera of the Baltic Amber. I. Ascalaphidae, Nymphidae, and Psychopsidae. *Psyche* 77:147-180.
- Młynarski, M., 1976. Testudines. *Encyclopedia of paleoherpetology*, Pt. 7. Stuttgart: Gustav Fischer Verlag, 130.
- Moore, R. C., Teichert, C., and Robison, R. A., eds., 1953-1982. *Treatise on invertebrate paleontology*. Lawrence, Kansas: Geol. Soc. Am. and Univ. Kansas Press.
- Morris, S. F., 1979. A new fossil terrestrial isopod with implications for the East African Miocene land fauna. *Br. Mus. Nat. Hist. Bull.* 32:71-75.
- Moy-Thomas, J. A., and Miles, R. S., 1971. *Paleozoic fishes*, 2nd ed. Philadelphia: Saunders, 259.
- Müller, A. H., 1963-1970. *Lehrbuch der Paläozoologie*. Jena: Veb Gustav Fischer Verlag.
- Mundel, P., 1979. The centipedes (Chilopoda) of the Mazon Creek. In Nitecki, M. H., (ed.), *Mazon Creek fossils*, New York: Academic Press, 361-368.
- Nelson, J. S., 1976. *Fishes of the world*. New York: Wiley-Interscience, 416.
- Niklas, K. J., Tiffney, B. H., and Knoll, A. H., 1985. Patterns in vascular land plant diversification: An analysis at the species level. In Valentine, J. W. (ed.), *Phanerozoic diversity patterns: Profiles in macroevolution*, Princeton: Princeton Univ. Press and Amer. Assoc. Adv. Sci. (this volume).
- Olsen, P. E., and Galton, P. M., 1977. Triassic-Jurassic tetrapod extinctions: Are they real? *Science* 197:983-986.
- Orlov, Yu. A., ed., 1958-1964. *Osnovy Paleontologii*. Moscow: Nedra.

- Padian, K., and Clemens, W. A., 1985. Terrestrial vertebrate diversity: Episodes and insights. In Valentine, J. W. (ed.), *Phanerozoic diversity patterns: Profiles in macroevolution*, Princeton, N.J.: Princeton Univ. Press and Amer. Assoc. Adv. Sci. (this volume).
- Piveteau, J., ed., 1952-1969. *Traité de Paléontologie*. Paris: Masson et Cie Éditeurs.
- Raup, D. M., Gould, S. J., Schopf, T. J. M., and Simberloff, D. S., 1973. Stochastic models of phylogeny and the evolution of diversity. *J. Geol.* 81: 525-542.
- Raup, D. M., and Sepkoski, J. J., Jr., 1982. Mass extinctions in the marine fossil record. *Science* 215:1501-1503.
- Rodendorf, B. B., ed., 1968. *Jurski Nasekomie Karatau*. Moscow: "Nauka," 252.
- Rolfe, W. D. I., Bonamo, P. M., Grierson, J. D. and Shear, W. A., 1983. The earliest land animals. *Am. Assoc. Adv. Sci.*, 149th Ann. Mtg., Abst. of Papers, 38-39.
- Romer, A. S., 1966. *Vertebrate paleontology*, 3rd ed. Chicago: Univ. Chicago Press, 468.
- Russell, D. A., 1975. Reptilian diversity and the Cretaceous-Tertiary transition in North America. *Geol. Assoc. Canada Spec. Pap.* (13):119-136.
- Schram, F. R., 1969. The stratigraphic distribution of the Paleozoic Eumalacostraca. *Fieldiana Geol.* 12:213-234.
- Schram, F. R., 1979. Worms of the Mississippian Bear Gulch Limestone of central Montana, USA. *San Diego Soc. Nat. Hist. Trans.* 19:107-120.
- Schram, F. R., and Schram, J. M., 1979. Some shrimp of the Madera Formation (Pennsylvanian) Manzanita Mountains, New Mexico. *J. Paleontol.* 53:169-174.
- Sepkoski, J. J., Jr., 1979. A kinetic model of Phanerozoic taxonomic diversity: II. Early Phanerozoic families and multiple equilibria. *Paleobiol.* 5: 222-251.
- Sepkoski, J. J., Jr., 1981. A factor analytic description of the Phanerozoic marine fossil record. *Paleobiol.* 7:36-53.
- Sepkoski, J. J., Jr., 1982a. Mass extinctions in the Phanerozoic oceans: A review. In Silver, L. T., and Schultz, P. H. (eds.), *Geological implications of impacts of large asteroids and comets on the earth*, *Geol. Soc. Amer. Spec. Pap.* 190.
- Sepkoski, J. J., Jr., 1982b. A compendium of fossil marine families. *Milwaukee Publ. Mus. Contr. Biol. Geol.* 51, 125.
- Simpson, G. G., 1953. *The major features of evolution*. New York: Columbia Univ. Press, 434.
- Solem, A., and Yochelson, E. L., 1979. North American Paleozoic land snails with a summary of other Paleozoic nonmarine snails. *U.S. Geol. Surv. Prof. Pap.* 1072, 42.
- Southcott, R. V., and Lange, R. T., 1971. Acarine and other microfossils from the Maslin Eocene, South Australia. *S. Aust. Mus. Rec.* 16(7):1-21.
- Steel, R., 1970. Saurischia. *Encyclopedia of paleoherpetology*, Pt. 14. Stuttgart: Gustav Fischer Verlag, 87.
- Steel, R., 1973. Crocodylia. *Encyclopedia of paleoherpetology*, Pt. 16. Stuttgart: Gustav Fischer Verlag, 116.
- Taylor, D. W., and Sohl, N. F., 1962. An outline of gastropod classification. *Malacologia* 1:7-32.
- Thompson, I., and Jones, D. S., 1980. A possible onychophoran from the middle Pennsylvanian Mazon Creek beds of northern Illinois. *J. Paleontol.* 54:588-596.
- Whalley, P. E. S., 1980. Neuroptera (Insecta) in amber from the Lower Cretaceous of Lebanon. *Br. Mus. Nat. Hist. Bull. (Geol.)* 33:157-164.
- Wighton, D. C., 1982. Middle Paleocene insect fossils from south-central Alberta. *Third N. Am. Paleont. Conv. Proc.* 2:577-578.
- Wilson, M. V. H., 1978. Paleogene insect faunas of western North America. *Quaest. Ent.* 14:13-34.
- Zangerl, R., 1981. Chondrichthyes. I. Paleozoic Elasmobranchii. *Handbook of Paleoichthyology*. Stuttgart: Gustav Fischer Verlag, 3A, 115.